

# Peach

## 2009 Pest Management Guide for Oregon

EM 8419-E

Revised February 2009

The chemicals, formulations, and rates listed for insect, mite, and disease control are among the best recommendations based on label directions, research, and orchard use experience. Only a thorough knowledge of the orchard, its variety, tree size and density, canopy characteristics, pest complex, and past pest problems will enable you to correctly select chemicals, rates, amount of water used per acre, and method of application for optimum pest control. Occasionally, different formulations of a product or like formulations containing a different amount of active ingredient also are registered and effective for use on the pests listed. These products also may be used; we do not intend to discriminate against them. You may wish to consult their labels and determine whether their use confers advantages over the products listed in this guide.

Always refer to the pesticide label for use instructions. It is the legal document regarding use patterns. Two questions frequently are asked about the chemical control of insects and diseases: "How much chemical do I use per acre?" and "What is the least amount of water I need per acre to apply in my concentrate sprayer?" Notice that the schedule below suggests an amount of formulated product (not active ingredient) to use per acre. This amount is based on a "typical" middle age and density orchard with moderate pest pressure. Common sense indicates that less material may be needed (less than that given) for 1- to 4-year-old orchards. Conversely, more chemical (within label limits) may be required for large, mature trees experiencing heavy pest pressure from multiple pests.

Many insecticide labels today indicate the minimum amount of water needed per acre to apply concentrate sprays of insecticides, as well as how to calculate the amount of chemical needed per acre in a concentrate sprayer. CHECK LABEL BEFORE SPRAYING!! Some label directions indicate dilute applications only. Also:

1. Make sure any tank-mixes of pesticides are compatible. For example, the elevated pH of some boron spray solutions weakens many insecticides. Boron also is incompatible with water-soluble packets.
2. Use adjuvants and spreader stickers with caution.
3. Heavy, brief rain or extended rain (0.75 inch for more than 24 hours) can remove pesticides from fruit and foliage. Reapplication may be necessary (within label limits).

### Important information

1. Be aware of worker protection standards (WPS). All new pesticide labels will provide orchard reentry intervals and personal protection equipment information.
2. Diazinon is now classified as a restricted-use pesticide due to bird toxicity. Maximum per-acre application rates have been reduced to 4 lb 50W, and the preharvest interval extended to 21 days.
3. *Orchard Pest Management, a Resource Book for the Pacific Northwest, 1993* (edited by Beers, Brunner, Willet, and Warner, published by Good Fruit Grower, Yakima, WA) has a comprehensive list of the tree fruit insect and mite pests, covering life histories, damage, detection, monitoring, and pest management. It is one of our primary sources of information in developing this pest management guide and the most complete reference on orchard use of the principles of integrated pest management.
4. Guthion has been cancelled.

### Stages

Dormant and delayed dormant

(stages 0, 1, and 2)

Prebloom (stage 3)

Popcorn (stages 4-5)

Full bloom (stage 7)



Not shown

Petal fall

Shuck split to shuck fall

Summer

Preharvest

Postharvest

Illustration courtesy of Washington State University Extension

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

## Peach Pest Control Recommendations

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

### Dormant and Delayed Dormant (just before buds open and before eggs hatch—Stages 0, 1, and 2)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
Cytospora canker	None	Remove and destroy dead cankered limbs.

#### Peach leaf curl and shothole

*Note:* Apply first leaf curl spray when 50% of the leaves have fallen and again at delayed dormant in late February before floral buds begin to open. An additional spray may be needed during the dormant season for shothole control depending on material selected. Adding a spreader sticker will increase the effectiveness of some of these leaf curl sprays. Even curl-resistant cultivars such as Frost need protection during the first few years. East of the Cascades, in southern Oregon, and in low-rainfall areas, a delayed dormant application alone should be effective.

bordeaux 12-12-100	—	—
Bravo Weather Stik	3-4 pt	Effective if used all dormant season long. Do not add a spreader sticker. 12-hour reentry.
Echo 720	3-4 pt	Effective if used all dormant season long. Do not add a spreader sticker. 12-hour reentry.
Kocide 2000	12 lb	Effective only on shothole if used during the dormant season. Many other copper products are labeled, such as C-O-C-S and Nordox. 24-hour reentry. Copper-Count-N has a 12-hour reentry.
lime sulfur (29%)	32 gal	A very effective product for leaf curl only. 48-hour reentry.
Nu-Cop 50DF	8-16 lb	Effective only on shothole if used during the dormant season. Many other copper products are labeled, such as C-O-C-S and Nordox. 24-hour reentry. Copper-Count-N has a 12-hour reentry.
Ziram 76DF	6-8 lb	A very effective product for both leaf curl and shothole. 48-hour reentry.

#### Scale, mite, and aphid eggs, peach twig borer

horticultural mineral oil (HMO) + diazinon 50WP	4-6 gal	When using a WP formulation with oil, fill sprayer tank one-third full with water, turn on agitator, slowly add the WP, fill tank one-half full with more water, add oil. Keep agitator running, finish filling. Do not exceed 1 application per season.
or Esteem 35WP	4-5 oz	<b>14-day PHI.</b>
or Success 2L	4-8 oz	<b>14-day PHI for peaches, 1-day PHI for nectarines.</b>

### Prebloom (prepink—Stage 3)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
------------------------------	-------------------------------	---

#### Shothole borer

*Note:* This pest has two and possibly three generations in the Willamette Valley.

endosulfan 50WP	1.5 lb/100 gal water	Determine approximate acreage infested; spray limbs and/or trunk to runoff. Spray infested limbs when new sawdust appears in existing shotholes (end of February or early March). Do not exceed 3 lb ai/A per year.
-----------------	----------------------	---

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

## Popcorn (just before petals begin to open—Stages 4-5)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
<b>Brown rot blossom blight</b>		
<i>Note:</i> See footnote 3, page 8.		
Abound	12-15.5 fl oz	See footnote 5, page 8. 4-hour reentry. <b>0-day PHI.</b>
Bravo Weather Stik	3-4.1 pt	Do not apply after shuck split. 12-hour reentry.
Bumper 41.8EC	4 oz	24-hour reentry. <b>0-day PHI.</b>
Captan 80WDG	2.5-5 lb	24-hour reentry
Elevate 50WDG	1-1.5 lb	12-hour reentry. <b>0-day PHI.</b>
Elite 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Indar 2F	6 fl oz	12-hour reentry. <b>0-day PHI.</b>
Orbit	4 oz	12-hour reentry. <b>0-day PHI.</b>
Orius 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Pristine	10.5-14.5 oz	See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
PropiMax EC	4 fl oz	Generic Orbit. 24-hour reentry. <b>0-day PHI.</b>
Quash	2.5-4 oz	12-hour reentry. <b>14-day PHI</b>
Quilt	14 fl oz	Abound + Orbit. See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
Rovral	1-2 pt	See footnote 3, page 8. 24-hour reentry.
Scala SC	9-18 fl oz	Do not exceed 3 applications alone. <b>2-day PHI.</b>
Thiram Granuflo	3.9-5.1 lb	24-hour reentry. <b>7-day PHI.</b>
Tilt	4 oz	12-hour reentry. <b>0-day PHI</b>
Topsin 4.5FL	20-30 oz	Tank-mix with another fungicide. 24-hour reentry. <b>1-day PHI.</b>
Vanguard 75WG	5 oz	Do not exceed 10 oz/A/season. Buffer to a pH of 5 to 7 if mixing with Rovral. 12-hour reentry.

## Peach twig borer, leafrollers, aphids, eyespotted bud moth, stinkbugs

*Note:* This is the most satisfactory time to apply green peach aphid-twig borer combination sprays. Asana, Pounce, and Ambush also are effective and registered for use, but may cause spider mite problems.

endosulfan 50WP	4 lb	Do not exceed 2 applications per year or 3 lb ai/A per year. If green peach aphid is a problem, use endosulfan. <b>30-day PHI.</b>
-----------------	------	--

## Peach twig borer, leafrollers, bud moth

<i>Bacillus thuringiensis</i>	1-2 lb	<i>Bt</i> products are stomach poisons. Complete coverage and 2 to 3 sprays usually are required for satisfactory control. Follow the label rates for individual products. <b>0-day PHI.</b>
Delegate	3-7 oz	<b>14-day PHI.</b>

## Silver or rust mites (if a problem)

endosulfan 50WP	2-3 lb	Do not exceed 2 applications per year or 3 lb ai/A per year. <b>30-day PHI.</b>
Nexter	10.67 oz	Do not exceed 2 applications per year. <b>7-day PHI.</b>
Vendex 50WP	1-2 lb	2-day reentry. Do not exceed 1.5 lb ai/A per year. <b>14-day PHI.</b>

## Thrips

*Note:* See footnote 4, page 8.

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

## Full bloom (Stage 7)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
------------------------------	-------------------------------	---

### Brown rot blossom blight

*Note:* See the popcorn stage for materials and rates to use.

## Petal Fall

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
------------------------------	-------------------------------	---

### Peach twig borer, leafrollers, oriental fruit moth, aphids

*Note:* Make petal fall spray if popcorn spray was missed or if orchard was heavily infested the previous season.

diazinon 50WP	4 lb	Do not exceed 1 application per season. <b>21-day PHI.</b>
Delegate	3-7 oz	<b>14-day PHI.</b>
Imidan 70WP	3.5-4.25 lb	<b>14-day PHI.</b>

### Peach twig borer, leafrollers, bud moth

<i>Bacillus thuringiensis</i>	1 lb	<i>Bt</i> products are stomach poisons. Complete coverage and 2 to 3 sprays usually are required for satisfactory control. <b>0-day PHI.</b>
Delegate	3-7 oz	<b>14-day PHI.</b>
Success 2L	4-8 oz	<b>14-day PHI.</b>

### Stink bugs

*Note:* Asana, Pounce, Ambush, and Guthion 50WP Solupak also are effective and registered for use, but may cause spider mite problems.

endosulfan 50WP	4 lb	Do not exceed 2 applications per year or 3 lb ai/A per year. <b>30-day PHI.</b>
-----------------	------	---

### Green peach aphid

Assail 30SG	2.5-5.3 oz	<b>7-day PHI.</b>
Provado 1.6F	4-8 oz	<b>0-day PHI.</b>

### Thrips

*Note:* See footnote 4, page 8.

### Brown rot blossom blight (for high rainfall areas)

*Note:* See the popcorn stage for list of materials and remarks.

## Shuck Split to Shuck Fall

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
------------------------------	-------------------------------	---

### Shothole (Coryneum blight)

Abound	11-15 fl oz	See footnote 5, page 8. 4-hour reentry. <b>0-day PHI.</b>
Bravo Weather Stik	3-4 pt	Do not apply past shuck split. 12-hour reentry.
Captan 80WDG	5 lb	24-hour reentry.
Echo 720	3-4 pt	Do not apply past shuck split. 12-hour reentry.
Gem	6-8 oz	Do not use with organosilicate surfactants. 12-hour reentry. <b>1-day PHI.</b>
Pristine	10.5-14.5 oz	See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
Ziram 76DF	6-8 lb	<b>30-day PHI.</b>

Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.

**CONTINUED—Shuck Split to Shuck Fall**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Powdery mildew</b>		
Abound	11-15.5 fl oz	See footnote 5, page 8. 4-hour reentry. <b>0-day PHI.</b>
Bumper 41.8EC	4 oz	24-hour reentry. <b>0-day PHI.</b>
Elite 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Gem	6-8 oz	Do not use with organosilicate surfactants. 12-hour reentry. <b>1-day PHI.</b>
Indar 2F	6 fl oz	12-hour reentry. <b>0-day PHI.</b>
Microthiol Disperss	10-20 lb	80% sulfur. Do not use a spreader sticker. 24-hour reentry.
Orbit	4 oz	12-hour reentry. <b>0-day PHI.</b>
Pristine	10.5-14.5 oz	See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
PropiMax EC	4 fl oz	24-hour reentry. <b>0-day PHI.</b>
Quilt	14 fl oz	Abound + Orbit. See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
Rally 40WSP	2.5-6 oz	24-hour reentry. <b>0-day PHI.</b>
Thiolux Jet	10-30 lb	80% sulfurs. 24-hour reentry.
Tilt	4 oz	12-hour reentry.

**Summer (first spray about 5 weeks before harvest or earlier if disease appears; repeat at weekly intervals; last application just before harvest)**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Powdery mildew (if found before pit hardening)</b>		
<i>Note:</i> See footnote 6, page 8.		
Bumper 41.8EC	4 oz	24-hour reentry. <b>0-day PHI.</b>
Elite 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Gem	6-8 oz	Do not use with organosilicate surfactants. 12-hour reentry. <b>1-day PHI.</b>
Indar 2F	6 fl oz	12-hour reentry. <b>0-day PHI.</b>
JMS Stylet oil	1-2 gal/ 100 gal water	Need good coverage when trees are dry. 4-hour reentry.
Orbit	4 oz	12-hour reentry. <b>0-day PHI.</b>
Pristine	10.5-14.5 oz	See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
PropiMax EC	4 fl oz	24-hour reentry. <b>0-day PHI.</b>
Rally 40WSP	2.5-6 oz	24-hour reentry. <b>0-day PHI.</b>
Tilt	4 oz	12-hour reentry. <b>0-day PHI.</b>

**Cucumber beetle**

Sevin XLR Plus	2-3 qt	<b>3-day PHI.</b>
----------------	--------	-------------------

**Shothole borer (if a problem)**

*Note:* Emergence in March to September, with three generations per year.

endosulfan 50WP	1.5 lb/ 100 gal water	Do not exceed 2 applications per year or 6 lb/A per year. Treat trunks and limbs. <b>30-day PHI.</b>
-----------------	--------------------------	--

**Spider mites**

M-Pede	1-2% solution	Potassium salts of fatty acids. <b>0-day PHI.</b>
--------	---------------	---

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

**CONTINUED—Summer**

<b>Pest or disease/ Material</b>	<b>Amount of product per acre</b>	<b>Comments/Reentry interval/Preharvest interval (PHI)</b>
<b>Spider mites, peach silver mite</b>		
Acaramite 50WS	0.75-1 lb	<b>3-day PHI.</b>
Apollo 50SC	4-8 fl oz	Make one application of Apollo or Savey. <b>21-day PHI.</b>
endosulfan 50WP	3-4 lb	Do not exceed 2 applications per season or 3 lb ai/A per season. Endosulfan is not effective on spider mites. <b>30-day PHI.</b>
Envidor	16-18 oz	<b>7-day PHI.</b>
Pyramite 60WSP	6.6-13.2 oz	Do not exceed 26.4 oz/A per season. <b>7-day PHI.</b>
Savey 50DF	3-6 fl oz	Will not control adults. Apply only once per season. <b>28-day PHI.</b>
Vendex 4L	1-2 pt	Do not exceed 2 applications per season or 1.5 lb ai/A per year. <b>14-day PHI.</b>
<b>Earwigs</b>		
Sevin 80S	0.6-0.9 lb/ 100 gal water	Apply around bases of trees and on trunks. Do not apply Sevin on cover crop blooms because of hazard to bees. <b>3-day PHI.</b>
<b>Peach twig borer, oriental fruit moth</b>		
<i>Note:</i> Apply twig borer and fruit moth sprays in early June or time sprays with pheromone traps. Apply cover spray about 14 days after pheromone traps average 2 moths per trap for first adult generation (May-June) or 5 moths per trap for second adult generation (July-August). Asana, Pounce, and Ambush also are registered but may cause spider mite problems.		
Altacor	3-4.5 oz	<b>10-day PHI.</b>
Assail 30SG	5.3-8 oz	<b>7-day PHI.</b>
Delegate	3-7 oz	<b>14-day PHI.</b>
diazinon 50WP	4 lb	Do not exceed 1 application per season. <b>21-day PHI.</b>
endosulfan 50WP	4-5 lb	Do not exceed 2 applications per year or 3 lb ai/A per year. <b>30-day PHI.</b>
Imidan 70WP	3.5-4.25 lb	<b>14-day PHI.</b>
Success 2L	4-8 fl oz	Do not exceed 29 oz/A per year. <b>14-day PHI.</b>
<b>San Jose scale, Lecanium scale crawlers (mid-June to early July)</b>		
diazinon 50WP	4 lb	Do not exceed 1 application per season. This spray is effective only on the crawler stages of scales. <b>21-day PHI.</b>
<b>Peach tree borer</b>		
<i>Note:</i> Timing usually is first or second week in July, and again 3 weeks later in August. If pheromone traps are used, place in orchard in June. Position traps about 2-3 feet from ground surface. Make first application 2 weeks after first consistent trap catches. Asana, Pounce, Ambush also registered for peach tree borers. Preharvest intervals are 14 days.		
endosulfan 50WP	—	Mix 1.5 lb/100 gal of water and spray trunk and lower limbs to runoff. Do not exceed 2 applications per year or 6 lb/A per year. <b>21-day PHI as trunk spray.</b>
Lorsban 4E	—	Mix 3 qt/100 gal of water and apply once per season as coarse low-pressure spray to trunks and lower crotches of peach trees. <b>14-day PHI.</b>

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Preharvest

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
<b>Brown rot fruit rot (if rain is forecasted)</b>		
Abound	12-15.5 fl oz	See footnote 5, page 8. 4-hour reentry. <b>0-day PHI.</b>
Bumper 41.8EC	4 oz	24-hour reentry. <b>0-day PHI.</b>
Captec 4L	4 qt	4-day reentry.
Elite 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Indar 2F	6 fl oz	12-hour reentry. <b>0-day PHI.</b>
Microthiol Disperss	10-20 lb	24-hour reentry.
Orbit	4 oz	12-hour reentry. <b>0-day PHI.</b>
Orius 45DF	4-8 oz	12-hour reentry. <b>0-day PHI.</b>
Pristine	10.5-14.5 oz	See footnote 5, page 8. 12-hour reentry. <b>0-day PHI.</b>
PropiMax EC	4 fl oz	24-hour reentry. <b>0-day PHI.</b>
Quash	2.5-4 oz	12-hour reentry. <b>14-day PHI</b>
Quilt	14 fl oz	Abound + Orbit. See footnote 5, page 8. 24-hour reentry. <b>0-day PHI.</b>
Scala SC	9-18 fl oz	Do not exceed 3 applications alone. <b>2-day PHI.</b>
Thiram Granuflo	3.9-5.1 lb	24-hour reentry. <b>7-day PHI.</b>
Tilt	4 oz	12-hour reentry. <b>0-day PHI.</b>
Topsin 4.5FL	20-30 oz	12-hour reentry. Tank-mix with another fungicide. <b>1-day PHI.</b>

## Postharvest (September-October)

Pest or disease/ Material	Amount of product per acre	Comments/Reentry interval/Preharvest interval (PHI)
Cytospora canker	None	Paint trunks with whitewash to help prevent winter injury.

### Peach leaf curl and shothole

*Note:* Apply first leaf curl spray when 50% of leaves have fallen and again at delayed dormant in late February before floral buds begin to open. An additional spray may be needed in dormant season for shothole control depending on material selected. Adding a spreader sticker increases effectiveness of some leaf curl sprays. Even curl-resistant cultivars such as Frost need protection during the first few years.

bordeaux 12-12-100	—	—
Bravo Weather Stik	3-4 pt	Effective if used all dormant season long. Do not add a spreader sticker. 12-hour reentry.
Echo 720	3-4 pt	Effective if used all dormant season long. Do not add a spreader sticker. 12-hour reentry.
Kocide 2000	12 lb	Effective only on shothole if used during the dormant season. Many other copper products are labeled, such as C-O-C-S and Nordox. 24-hour reentry. Copper-Count-N has a 12-hour reentry.
lime sulfur (29%)	32 gal	A very effective product for leaf curl only. 48-hour reentry.
Nu-Cop 50DF	8-16 lb	Effective only on shothole if used during the dormant season. Many other copper products are labeled, such as C-O-C-S and Nordox. 24-hour reentry. Copper-Count-N has a 12-hour reentry.
Ziram 76DF	6-8 lb	A very effective product for both leaf curl and shothole. 48-hour reentry.

### Shothole borer (third generation, postharvest control)

*Note:* Determine approximate acreage infested and spray limbs and/or trunk to runoff. Spray infested limbs when new sawdust appears in existing shotholes, September/ October.

endosulfan 50WP	1.5 lb/100 gal water	Do not exceed 2 applications per year or 6 lb/A per year.
Lorsban 4E	3 qt/100 gal water	Apply as a trunk spray. Do not exceed 1 application per season.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Footnotes

1. Bordeaux mixture may be combined with horticultural mineral oil for both peach leaf curl and Lecanium scale control. No sticker is needed with this combination.
2. Bordeaux 12-12-100 means 12 pounds of copper sulfate plus 12 pounds of lime in 100 gallons of water. In any bordeaux formula, the ingredients always are listed in the same order—copper sulfate, hydrated lime, then gallons of water.
3. Fungal pathogens have shown resistance to many fungicides when one is used exclusively. Alternate or tank-mix fungicides with different modes of action. Fungicides from different groups have different modes of action.
4. Surface scarring on stone fruits can be the result of many factors. Certain insects such as thrips deform and bronze flower buds and blossoms. Thrips can scar fruit by feeding on or laying eggs in the fruit. Most significant damage usually is during and shortly after pollination.

Lygus and stink bugs also damage stone fruit at this time. Buds are injured, flowers can be sterile, and fruit may be dimpled, distorted, and “pock-marked.”

Damage from the above pests is sporadic and is only occasional in some Valley orchards. Sometimes only portions of orchards or border rows are damaged. Best timing to prevent damage also coincides with pollination periods. Even though some varieties may be wind pollinated, bees can boost yield, often are present, and must be protected. Prebloom and petal fall sprays of endosulfan (Lygus and stink bugs), spinosad (Success), or Carzol (thrips) should be applied in the evening after bee activity.

**Be sure fruit scarring is the result of insects before applying these sprays.**

5. Do not use more than two consecutive applications before switching to another fungicide in a different family or group with a different mode of action. Sprayers used for Abound should **not be used on apples** such as Gala, Cox’s Orange Pippin, and McIntosh.
6. Powdery mildew may be a problem in some years. Nearby roses are an alternate host for this fungus. Scout for first occurrence. Chemical control is not needed after pit hardening. A similar disease called rusty spot comes from local apple trees with powdery mildew.

---

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Table 1. Effectiveness of Fungicides and Bactericides for Peach Disease Management\*

Fungicide	Fungicide group #	Brown rot (blossom blight)	Brown rot (fruit rot)	Peach leaf curl	Powdery mildew	Shothole	Pseudomonas bacterial canker
Abound	11	(Fair to) good**	(Fair to) good**	??	Excellent**	Fair-good	Not effective
Botran	14	Fair	Fair	Slight	Not effective	??	Not effective
Bravo	M5	Good-fair	Not registered	Good	Not registered	Good	Not effective
Bumper	3	Excellent	Excellent	Slight	Excellent**	Slight	Not effective
Captan	M4	Good	Fair-good	Slight	Not effective	Fair-good	Not effective
Copper-based products	M1	Slight	Not registered	Fair-good	Slight	Good	Fair-excellent**
Echo	M5	Good-fair	Not registered	Good	Not registered	Good	Not effective
Elevate	17	Good-excellent	Good-excellent	??	Not effective	??	Not effective
Elite	3	Good-excellent	Good-excellent	Fair	Excellent**	??	Not effective
Ferbam	M3	Fair	Not registered	Good	Not registered	Good	Not effective
Gem	11	Fair-good**	Fair-good	??	Excellent**	??	Not effective
Indar	3	Excellent**	Excellent**	Fair	Excellent**	??	Not effective
JMS Stylet oil	Not classified	??	??	??	Good	??	??
lime sulfur	M2	Not recommended	Not recommended	Excellent	Not recommended	Slight	Not effective
Orbit	3	Excellent	Excellent	Slight	Excellent**	Slight	Not effective
Orius	3	Good-excellent	Good-excellent	Fair	Excellent**	??	Not effective
Pristine	11+7	Good-excellent	Good-excellent	??	Excellent	??	Not effective
PropiMax	3	Excellent	Excellent	Slight	Excellent**	Slight	Not effective
Quash	3	Good to excellent**	Good to excellent**	??	Suspect good**	??	Not effective
Quilt	11+3	Excellent	Excellent	Slight	Excellent**	Fair-good	Not effective
Rally	3	Good	Good	??	Excellent**	Slight	Not effective
Rovral	2	Good**	Not registered	Slight	Not effective	Fair-good	Not effective
Scala	9	Good	Good	??	None	??	None
Sulfur	M2	Fair	Fair (good)	Slight	Good	Not effective	Not effective
Syllit	M7	??	Not registered	??	Not registered	??	None-slight
Thiram	M3	Good	Good	Good	Not effective	??	Not effective
Tilt	3	Excellent	Excellent	Slight	Excellent**	Slight	Not effective
Topsin	1	Good**	Good**	Not effective	Good**	Not effective	Not effective
Vanguard	9	Good**	Not registered	??	Not effective	Fair	??
Ziram	M3	Fair	Not registered	Excellent	Not effective	Good-excellent	Not effective

\*These ratings are relative rankings based on labeled application rates, good spray coverage, and proper spray timing. Actual levels of disease control will be influenced by these factors in addition to cultivar susceptibility, disease pressure, and weather conditions.

\*\*Resistant pathogens will lower the effectiveness of these fungicides.

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Table 2. Registered herbicides for orchard crops in Oregon

This abbreviated table was prepared by Jeff Olsen, OSU Extension horticulturist, and Ed Peachey, OSU horticultural crop weed scientist. For a more complete guide to orchard crop weed management, please refer to the *PNW Weed Management Handbook*, available in print or online ([http://pnwpest.org/pnw/weeds?21W\\_HORT01.dat](http://pnwpest.org/pnw/weeds?21W_HORT01.dat)). For access to herbicide labels, visit the CDMS Agro-chemical database (<http://www.cdms.net/LabelsMsds/LMDefault.aspx>) or Greenbook (<http://www.greenbook.net/>).

### Site preparation

Material	Uses	Rate
glyphosate (numerous product names)	—	Read label carefully for crops listed and geographic location.

### New plantings (newly planted trees less than 1 year old), nonbearing

Material	Uses	Rate
napropamide (Devrinol)	All except chestnuts	5 lb ai/A (50 lb/A Devrinol 10-G)
oryzalin (Surflan)	—	2-6 lb ai/A (2-6 qt/A Surflan AS)
trifluralin (Treflan 4EC)	Newly established, nonbearing and bearing, except pears, apples, cherries, and hazelnuts	0.5-1 lb ai/A (1-2 pt/A Treflan 4EC)
isoxaben (Gallery or Gallery T&V)	Nonbearing crop only, except chestnuts	0.495-0.998 lb ai/A (0.66-1.33 lb/A product)
isoxaben + trifluralin (Snapshot 2.5TG)	Nonbearing crop only, except chestnuts	5 lb ai/A (200 lb/A Snapshot 2.5TG)
trifluralin + isoxaben + oxyfluorfen (Showcase)	Nonbearing crops only, including apricots, cherries, nectarines, peaches, plums, and prunes; excluding apples, pears, hazelnuts, walnuts, and chestnuts	2.5-5 lb ai/A (100-200 lb/A Showcase)
pronamide (Kerb)	Nonbearing fruit trees	1-4 lb ai/A (2-8 lb/A). Rate depends on species present and soil texture.
fluazifop (Fusilade DX)	—	0.25-0.375 lb ai/A (16-24 oz/A Fusilade DX). Refer to specific grassy weeds on label.
sethoxydim (Poast)	—	0.28-0.47 lb ai/A (1.5-2.5 pt/A product)
clethodim (Envoy, Prism, Select)	Nonbearing trees only	0.06-0.125 lb ai/A (6-8 oz/A Select)
paraquat (Gramoxone, Inteon, Firestorm)	—	0.625-1 lb cation/A (2.5-4 pt/A Gramoxone; 1.7-2.7 pt/A Firestorm)
glyphosate (numerous product names)	—	Read label carefully for crops listed and geographic location.
glyphosate (numerous product names)	—	Wiper: 33% solution
glufosinate ammonium (Rely)	Apples, hazelnuts, and walnuts	0.75-1.5 lb ai/A (3-6 qt/A Rely)
oxyfluorfen (Goal 2XL)	—	1.25-2 lb ai/A (5-8 pt/A Goal 2XL)

Continued on next page

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Established plantings/Applications that persist in the soil

Material	Uses	Rate
pendimethalin (Prowl H <sub>2</sub> O)	Bearing fruits and nuts (H <sub>2</sub> O formulation); 3.3EC formulation for nonbearing only	Prowl H <sub>2</sub> O: 1.9-3.8 lb ai/A (2-4 qt/A), depending on desired length of weed control. Prowl 3.3EC: 1.98-3.96 lb ai/A (2.4-4.8 qt/A)
flumioxazin (Chateau WDG)	Nut trees only	0.188-0.38 lb ai/A Chateau WDG. Refer to label for organic matter, soil types, and rates for various broadleaf weeds.
flumioxazin (Chateau SW)	Pome and stone fruit trees	Preemergence: 0.38 lb ai/A (12 oz/A Chateau SW). Postemergence: 0.188-0.38 lb ai/A (6-12 oz/A Chateau SW). See label for organic matter, soil types, and rates for various broadleaf weeds.
dichlobenil (Casoron CS)	Apples, pears, cherries, and hazelnuts only, bearing and nonbearing	1.96-3.92 lb ai/A (1.4-2.8 gal/A Casoron CS)
dichlobenil (Casoron)	Apples, pears, cherries, and hazelnuts	4-6 lb ai/A (100-150 lb/A Casoron)
diuron (Karmex DF and others)	Apples, pears, peaches, hazelnuts, and walnuts only	See label for crop-specific application rates.
simazine (Princep Caliber 90 is a Special Local Needs label for sweet cherries only). Tart cherries have a federal label for Princep 4L, Caliber 90, and other simazine herbicides in Oregon.	Except chestnuts, apricots, prunes, and nectarines	See product labels for rates.
terbacil (Sinbar 80)	Bearing apples and peaches only; labeled for nonbearing apricot, pear, apple, cherry, peach, and plum	0.4-0.8 lb ai/A (0.5-1 lb/A), newly established; 2-4 lb/A Sinbar depending on soil type
norflurazon (Solicam)	Except chestnuts	1.97-7.8 lb ai/A (2.5-10 lb/A Solicam)
napropamide (Devrinol)	Except chestnuts	5 lb ai/A (50 lb/A Devrinol 10G)
oryzalin (Surflan AS)	Except apples, cherries, nectarines, peaches, prunes, and walnuts	2-6 lb ai/A (2-6 qt/A Surflan)
pronamide (Kerb)	Except apricots, hazelnuts, and walnuts	1-4 lb ai/A (2-8 lb/A Kerb 50W)
trifluralin (Terflan 4EC)	Except apples, pears, cherries, and hazelnuts	0.5-1 lb ai/A (1-2 pt/A Terflan 4EC)
oxyfluorfen (Goal 2XL)	—	1.25-2 lb ai/A (5-8 pt/A Goal 2XL)
pendimethalin	Supplemental labels for Prowl H <sub>2</sub> O on bearing stone and pome fruits and nuts	1.9-3.8 ai/A (2-4 qt/A Prowl H <sub>2</sub> O), depending on desired length of control

Continued on next page

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

## Established plantings/Postemergence contact and translocated herbicides

Material	Uses	Rate
paraquat (Gramoxone Max, Firestorm)	—	0.625-1 lb cation/A (2.5-4 pt/A Gramoxone; 1.7-2.7 pt/A Firestorm)
oxyfluorfen (Goal 2XL)	—	1.25-2 lb ai/A (5-8 pt/A Goal 2XL)
glyphosate (numerous product names)	—	Spray: read label carefully for crops listed and geographic location.
glyphosate (numerous product names)	—	Wiper: 33% solution
glufosinate ammonium (Rely)	Apples, hazelnuts, and walnuts	0.75-1.5 lb ai/A (3-6 qt/A Rely)
2,4-D amine (Saber, Weed-Rhap A4D, Dri-Clean Herbicide)	—	0.95-1.4 lb ae/A (2-3 pt/A Saber)
clopyralid (Stinger)	Stone fruits only	0.12-0.25 lb ae/A (0.33-0.66 pt/A Stinger)
rimsulfuron (Matrix FNV)	All fruit and nut crops	0.063 lb ai/A (4 oz/A Matrix FNV)
halosulfuron (Sanda)	Nut crops only	0.032-0.062 lb ai/A (0.66-1.33 oz/A)
fluzifop (Fusilade DX)	Except bearing apples, pears, chestnuts, and walnuts; labeled for nonbearing pears	0.25-0.375 lb ai/A (16-24 oz/A Fusilade DX). Refer to specific grassy weeds listed on label.
sethoxydim (Poast)	Except prunes and plums	0.28-0.47 lb ai/A (1.5-2.5 pt/A Poast)

## Grass suppression in row middles (chemical mowing)

Material	Uses	Rate
glyphosate (numerous product names)	—	Read label carefully for crops listed and geographic location.
sethoxydim (Poast)	—	0.09 lb ai/A (0.5 pt/A Poast)

## OSU Internet resources for plant protection

Information regarding plant protection is available from several sources at OSU. The following listings are excellent examples:

- OSU Integrated Plant Protection Center. Online weather data and degree day information for insect pests and diseases (<http://pnwpest.org/wea/index.html>)
- Codling moth development information (<http://pnwpest.org/cgi-bin/ddmodel.pl?clm>)
- Apple scab infection season information (<http://pnwpest.org/cgi-bin/ddmodel.pl?spp=asc>)
- Pear scab infection season information (<http://pnwpest.org/hr/>)
- Pear scab infection period information for the Hood River Valley (<http://pnwpest.org/hr/>)
- Fire blight risk information (<http://pnwpest.org/cgi-bin/ddmodel.pl?fbl>)  
Directions for the use of each model are available at each site.
- OSU Botany and Plant Pathology Department. Site of “Online Guide to Plant Disease Control” (<http://ipmnet.org/plant-disease/>). Disease symptom descriptions, pictures of disease symptoms, and other information helpful in plant protection
- Pacific Northwest Insect Management Handbook (<http://pnwpest.org/pnw/insects>)
- Pacific Northwest Weed Management Handbook (<http://pnwpest.org/pnw/weeds>)

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*

# Basic Elements of Safe Pesticide Use

- Always read the label with care. This is the first step in selecting the right material for the job. Never rely on your memory. Before opening the container, pay strict attention to warnings and cautions printed on the label.
- Keep all pesticide and spray materials out of the reach of children, pets, and irresponsible persons. Storage outside of the home, away from food and feed, and under lock and key is the safest method.
- Store only in the original container and keep tightly closed.
- NEVER smoke, eat, or drink while applying pesticides.
- Avoid inhalation or direct contact. Always wear protective clothing and safety devices as recommended on the label.
- Avoid spills. If spills occur, take immediate action to remove contaminated clothing and wash thoroughly.
- After each application, bathe and change to clean clothing. Wash clothing after each use. Always use fresh clothing when starting new application.
- Avoid contamination of fish ponds and water supplies. Cover feed and water containers when treating around livestock or pet areas.
- Keep separate equipment for use with hormone-type herbicides to avoid accidental injury to susceptible plants. Also avoid applications under wind conditions that could create drift to nontarget areas.
- Rinse empty containers three times before disposing of them. Add the rinse to the spray tank and dispose of containers according to local regulations to avoid hazard to humans, animals, and the environment.
- Follow label directions for mixing and application to keep residues within the limits prescribed by law.
- Plan ahead. Discuss with your physician the materials you will be using during the season so that he or she can be prepared to provide the appropriate treatment in case of accidental exposure. If symptoms of illness occur, call the physician or get the patient to a hospital immediately. Always provide the medical personnel with as much information as possible.
- Be cautious when you apply pesticides. Know your legal responsibility as a pesticide applicator. You may be liable for injury or damage resulting from pesticide use.

## Oregon Poison Center

The Oregon Health Sciences University

3181 S.W. Sam Jackson Park Road, Room CB 550

Portland, OR 97201

Phone: 503-494-8968; Oregon Toll Free: 1-800-452-7165; Nationwide: 1-800-222-1222

**If a person has collapsed or is not breathing, dial 911.**

---

© 2009 Oregon State University. This publication may be photocopied or reprinted **in its entirety** for noncommercial purposes.

Prepared by Jeff Olsen, Extension horticulturist, Yamhill County, and Jay W. Pscheidt, Extension plant pathologist, Oregon State University. The information in this pest management guide is valid for 2009. The mention of commercial products in this publication does not constitute endorsement by the Oregon State University Extension Service, nor should exclusion be interpreted as criticism of any item, form, or service. Due to constantly changing laws and regulations, the Oregon State University Extension Service can assume no liability for the suggested use of chemicals contained in this guide. Pesticides should be applied according to the label directions on the pesticide container.

This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

Revised February 2008; revised February 2009

---

*Use only one material except where a combination is indicated. Follow label precautions when tank-mixing oils, fungicides, and insecticides. Materials are not listed in order of preference.*