

Extension Forestry

We're here to serve you!

<http://extension.oregonstate.edu/douglas/forestry>

OSU
Oregon State
UNIVERSITY

DOUGLAS
COUNTY OSU
EXTENSION
SERVICE

The OSU Extension Service has been your "front door" to Oregon State University since 1868.

PO Box 1165
1134 SE Douglas Ave
Roseburg, OR 97470

Phone: 541-672-4461
Fax: 541-672-4453



John Punches
Forestry Agent

john.punches@oregonstate.edu

Characteristics of Tree Species Common in Douglas County

Compiled by
John Punches



Douglas-fir (*Pseudotsuga menziesii*)

- Most common west coast tree
- Fast growing on good sites, likes mineral soil
- Does not tolerate extreme wet or dry conditions
- Shade intolerant
- Often forms dense pure stands following fire
- Well suited to Douglas County's north and east facing slopes
- Oldest authentically dated tree was 1400 years old (500 years is more common)
- Growth extremely limited after 200 years
- May form tap root in deep soil, but more commonly stays near the surface
- Low root pressure
- Root system about diameter of crown
- Difficult to achieve natural regeneration after harvest
- Tallest specimen 330 feet.
- 250 feet height and 6 feet diameter common in old forests.

Pacific Silver Fir (*Abies amabilis*)

- Gray trunk and rigid, symmetrical crown with branches perpendicular to crown
- Maritime climate
- Cool summers
- 38 to 262 inches precipitation
- Must have adequate soil moisture
- Wide range of soils, prefers well-drained
- Found at high elevation in Douglas County (3500 to 6000 ft)
- Commonly associates with hemlock
- Can produce epicormic branches
- Can exceed 200 ft height and 24 inches in dbh
- Maximum reported age is 590 years
- May develop taproot
- Flat, shallow root system on poorly drained or shallow soils
- One of the most shade tolerant species in the Northwest

White Fir (*Abies concolor*)

- Rogue/Umpqua Divide, high elevations
- Wide variety of soils and soil conditions
- Growth best on moderately deep, well drained sandy-loam and clay-loam soils
- Frost sensitive
- Dry sites
- Very shade tolerant, but less so than western hemlock and perhaps western redcedar
- Grows best in full sunlight
- Adaptable rooting habit
- May reach 180 feet tall and 65 inches dbh

Grand Fir (*Abies grandis*)

- Rapid growing
- Has reached 231 ft height and 79 inches dbh
- Wide range in western Oregon and Washington
- Grows well on wide variety of soils
- Prefers deep, rich alluvial soils of valley bottoms
- May produce taproot
- Lives 250 to 300 years maximum
- Shade tolerant - more so than Douglas-fir, less so than hemlock
- May reach 250 ft in height and 60 inches in dbh

Subalpine Fir (*Abies lasiocarpa*)

- Smallest fir species
- Long, narrow conical crown
- Grows in cool, wet climate
- High elevations in Oregon
- Tolerates both wet and dry sites
- May reach 100 ft height and 24 inches dbh
- May exceed 250 years age
- Very shade tolerant
- Wide range of soils
- Generally shallow rooted
- Susceptible to windthrow

California Red Fir (*Abies magnifica*)

- Rogue/Umpqua divide, Crater Lake, Diamond Lake
- Limited to high elevations (5300 to 5900 ft)
- Cool/cold and moist climates
- Often found on young soils
- High stand densities
- Adaptable rooting habit
- Relatively shade tolerant
- Grows best in full sunlight
- Reaches 200 ft in height and 5 ft in diameter

Noble Fir (*Abies procera*)

- Crater Lake, Diamond Lake
- 200 ft tall and 5 ft in dbh
- Prefers moist, maritime climate
- Grows primarily at higher elevations (3500 to 5500 ft)
- Tolerates wide range of soils
- Must have adequate soil moisture
- Slow initial growth
- Commonly reach 45 to 60 inches dbh and 130 to 175 ft tall
- Largest is 278 ft tall and 108 inches dbh
- Frequently found in mixed stands
- Moderately windfirm
- Intermediately shade tolerant - least shade tolerant of any American true fir

Western Hemlock (*Tsuga heterophylla*)

- Humid areas of Pacific coast & northern Rockies
- Thrives in mild, humid climate with frequent fog and precipitation during growing season
- Confined to north aspects, stream bottoms, and seepage sites in dryer climates
- Grows on variety of soils, but growth decreases on compacted or clayey soils
- Does not tolerate surface water - needs 6 inches to 3 feet of drained soil
- Very shade tolerant
- Shallow rooted
- Very susceptible to fire and windthrow
- Sunscald possible in thinnings
- Grows in dense stands (up to 10,000 stems/acre)
- May reach 700 years age, but trees over 300 virtually stop growing
- Commonly reach 200 ft height and 40 inch dbh (max = 259 ft and 108 inch dbh)

Yew (*Taxus brevifolia*)

- Small shade-loving conifer often found in mature conifer stands
- Shade tolerant
- Usually an understory tree
- Slow growing
- May reach 24 inches dbh and 50 ft height
- Has hard, heavy, decay-resistant wood
- Prefers deep, moist, rocky or gravelly soils
- Dioecious (separate male and female flowering plants)

Redwood (*Sequoia sempervirens*)

- Mild, very humid climate
- Coastal region of northern California and far southern corner of Oregon
- Prefers fog
- Deep sandy loam or gravelly loam soils, likes flood plains
- Will not tolerate high levels of sodium or magnesium in soil
- Will reproduce vegetatively
- World's tallest tree (200 to 300 ft common, record is over 360 ft)
- Can reach over 240 inches diameter (dbh)
- No taproot, lateral roots large and wide-spreading
- Usually a dominant tree, yet very tolerant of shade
- Grows in very dense stands

Giant Sequoia (*Sequoiadendron giganteum*)

- Natural range is west slope of Sierra Nevada in central California
- Humid climate with dry summers
- Temperatures below -12 degrees F. limit range
- Deep, well-drained sandy loam soils
- Will grow on shallow, rocky soils if subsoil moisture is available
- 4500 to 6500 feet elevation
- Natural reproduction is tenuous - requires very specific conditions
- Long-lived and massive
- Commonly reach 250 ft in height and 120 to 240 inches dbh
- Many individuals 2000 to 3000 years old
- Forms significant taproot
- Shade intolerant

Port-Orford-Cedar (*Chamaecyparis lawsoniana*)

- Not a true cedar
- Range includes southwest Douglas County
- Cool wet winters and hot dry summers
- Moderate to high rainfall, fog common
- Grows on sand dunes, bogs, intermittent stream margins, and drier sites
- Best growth on deep soils from weathered sedimentary rock
- Prefers concave or sheltered slopes with seepage
- Relatively slow growing
- May exceed 560 years age
- May grow to 160 ft height and 40 inch dbh
- Shade tolerant
- Dense stocking common

Western Redcedar (*Thuja plicata*)

- Not a true cedar
- Prefers boggy habitats
- Can tolerate sites too dry for hemlock
- 28 to 260 inches of rain
- Susceptible to frost damage and windthrow
- Wide range of soils
- Tolerates low soil fertility but responds very well to fertile sites

Western Redcedar (*Thuja plicata*) (continued)

- Raises soil pH
- Prefers disturbed mineral soil seedbed
- Extensive rooting without taproot
- Will reproduce vegetatively
- Establishes in dense stands (1000/acre)
- Very shade tolerant (but less than hemlock)
- Found as high as 7500 ft in Oregon
- 200 ft tall and 10 feet or more in diameter
- Commonly lives 800 to 1000 years (2000 rare)

Incense-cedar (*Calocedrus decurrens*)

- Not a true cedar
- Slow growing
- Drought tolerant
- More shade tolerant than Douglas-fir
- Tolerates shade as seedling
- Prefers open sunlight as pole or mature
- Spreading and extensive root system
- Susceptible to root rots, especially annosus
- Grows on variety of soils
- May exceed 500 years age
- May reach 225 ft height and 150 inch dbh

Sitka Spruce (*Picea sitchensis*)

- Largest of world's spruces
- Grows in narrow strip along Pacific coast
- Requires humid air, likes summer fogs
- Best on deep, moist, well-aerated soils
- Growth poor on swampy sites
- Requires relatively high amounts of Ca, Mg, P
- May live 700 to 800 years
- Has reached 280 ft in height and over 16 ft dbh (current record tree is near Seaside OR)
- Shade tolerance between Doug-fir & hemlock
- Branches epicormically when opened to sunlight
- Susceptible to windthrow
- Great variability in rooting habit

Ponderosa Pine (*Pinus ponderosa*)

- Prefers coarse-textured soils
- Tolerates wide range of climate and elevations
- Tolerates low soil fertility
- Often found in groups of similar-aged trees rather than true uneven-aged structure
- Fire resistant when mature
- 300 to 600 year lifespan
- Vigorous taproot (3 to 6 feet)
- Intolerant to shade
- Holds 2 to 3 years needles, so crown has tufted appearance

Jeffrey Pine (*Pinus jeffreyi*)

- Primarily a California species, reaches into southwest corner of Douglas County
- Similar in appearance and wood characteristics to ponderosa pine

Jeffrey Pine (*Pinus jeffreyi*) (continued)

- Very cold hardy
- Drought hardy
- Adapted to short growing seasons
- Tolerates shallow, infertile, loamy to clayey & serpentine soils
- Will grow on any well-drained soil
- Widely associating with incense-cedar
- Grows more slowly than ponderosa pine during sapling stage, but more rapidly during pole stage
- May live 400 to 500 years
- May reach 200 ft height and 90 inch dbh
- Very windfirm
- Shade intolerant
- Holds needles for 5 to 8 years, so has more full crown than ponderosa pine

Lodgepole Pine (*Pinus contorta*)

- 4 varieties, we have 2 in Oregon: shore pine (var. *contorta*) and Rocky Mountain lodgepole (var. *latifolia*)
- Grows under wide variety of climatic conditions
- Relatively resistant to frost injury
- Very intolerant of shade, best in full sunlight
- Wide variety of soils, usually moist
- Tolerates bogs as well as dry, sandy sites
- Tolerates extremely infertile soils
- Prolific seed producer
- Cones are persistent and often serotinous
- Non-serotinous cones are prevalent in Oregon
- Regenerates in dense stands following fire
- May live 300 to 400 years, but this is rare
- May reach 20 inches dbh and 100 ft tall
- Taproots common in deep soil
- Shallow rooting common where hardpan or water table is encountered

Sugar Pine (*Pinus lambertiana*)

- Tallest and largest of the pines
- Reaches 200 ft in height and 60 inches dbh
- Occasionally exceed 500 years age
- Found at 1100 to 5400 ft elevation
- Temperature and precipitation vary widely
- Wide range of soil conditions
- Best on well-drained, acidic soils
- Often found on steep and rugged terrain
- Develops deep taproot
- More shade tolerant than ponderosa pine, but less so than incense-cedar and Douglas-fir

Western White Pine (*Pinus monticola*)

- Found in Cascade Range
- Commonly 190 ft tall and 70 inches dbh
- Has reached 239 ft in height and 78 inches dbh
- Maritime climates, wet winters & dry summers
- Tolerates wide range of soils
- Prefers deep porous soils

Western White Pine (*Pinus monticola*) (continued)

- Most common on poor, sandy soils
- Reaches 300 to 400 years old, rarely 500 years
- Root habit depends on soil conditions
- Roots tolerate dense soil
- Intermediate in shade tolerance
- Achieves dominant position in stand only following wildfire or clearcutting

Hardwoods

Red Alder (*Alnus rubra*)

- Most common hardwood in Pacific Northwest
- Lowland species
- Usually found within 125 miles of ocean and at elevations below 2400 feet
- Humid climates
- Needs 25 inches rain per year or access to ground water
- Tolerates wide range of soils
- Tolerates poor drainage and some flooding during growing season
- Prevalent along streams and in swamps or marshes
- Will sprout vigorously from stump when young
- Rapid juvenile growth
- Extensive, fibrous root systems
- Intolerant of shade - only black cottonwood requires more sunlight
- May experience epicormic branching following thinning
- Short-lived, maximum age is about 100 years
- Trees over 35 inches dbh are rare
- May reach 130 ft height on good site

Bigleaf Maple (*Acer macrophyllum*)

- Leaves up to 24 inches across
- Wide range of temperature and moisture conditions
- Often near streams and springs but also found on dry sites
- Prefers well-drained alluvial soils - growth is poor on shallow, rocky soils
- Sprouts prolifically after being cut
- Rapid juvenile growth
- Generally 50 ft tall and 20 inches dbh
- Has been know to exceed 100 ft in height and 130 inches dbh
- Maximum age appears to be about 200 years

Black Cottonwood (*Populus trichocarpa*)

- Largest of the American poplars
- Largest known hardwood tree in Western North America
- Wide range throughout Pacific Northwest
- Best growth in humid coastal forest, but tolerates dry climates
- Grows on variety of soils
- High soil acidity may restrict establishment
- Found at up to 5000 ft elevation in Cascade Range
- Forms extensive pure stands on alluvial soils
- Sprouts readily from cut stumps

Black Cottonwood (*Populus trichocarpa*) (continued)

- Cuttings easily rooted
- May attain pulpwood size in 10 to 15 years
- Grows well for as long as 200 years
- Exceptional trees have attained 120 inches dbh and 200 ft height
- Very intolerant of shade - grows best in full sunlight

Oregon Ash (*Fraxinus latifolia*)

- Common in riparian habitats
- May reach 250 years of age
- Fast growing during juvenile period, slow growing later
- Prefers cool summers and mild winters
- Occasionally found as high as 5000 ft elevation, usually not higher than 3000 ft
- Prefers poorly drained, moist bottom land with deep soil rich in organic matter
- May reach 80 ft in height and 30 inches dbh
- Generally shallow rooted
- Intermediate shade tolerance
- Self pruning is rapid where side shade is present

Willow (*Salix* spp.)

- Wide variety of species common in wet sites
- No timber value in Douglas County
- Used in basket weaving and other crafts
- Generally small and bushy in appearance

Oregon White Oak (*Quercus garryana*)

- Principal native oak in Oregon
- Diverse climates, ranging from cool and humid to hot and dry
- Grows on wide range of soils and sites
- Most competitive on poor sites
- Tolerates site conditions ranging from extremely dry to very moist
- One of few species that survive well on sites that cycle between extremely wet and extremely dry
- Sprouts abundantly from dormant buds following cutting or fire
- May reach 90 ft tall and 40 inches dbh on good sites
- Often stunted and of poor stem form on low quality sites
- Deep taproot and well-developed lateral root system
- Intermediate shade tolerance, but not tolerant to sustained overtopping by other trees

California Black Oak (*Quercus kelloggii*)

- Most common California oak, range stops just south of Eugene
- Does not grow near Pacific Ocean
- Climate typified by hot dry summers and cool moist winters
- Does not tolerate wet soils (so not commonly found where clay content is high)
- Found at 450 to 1000 ft elevation
- Commonly found mixed with Oregon White Oak and/or Madrone in Douglas County

California Black Oak (*Quercus kelloggii*) (continued)

- Sprouts profusely after trees are cut or burned
- Most commonly regenerated by sprouting following fire
- May live to be 500 years old
- Seldom exceeds 130 ft in height and 5 ft diameter
- Large spreading lateral roots and may form multiple tap roots
- Moderately shade tolerant in early life but becomes progressively intolerant to shade as it matures - will not tolerate overtopping for extended periods

Madrone (*Arbutus menziesii*)

- One of the most widely distributed tree species on the Pacific coast
- Broadleaved evergreen
- Prefers mild temperatures and prolonged cloudy periods
- Prefers wet, mild winters and cool, relatively dry summers with long frost-free periods
- Wide range of soils and terrains
- Generally not found above 3000 ft elevation in Douglas County
- Associates with wide range of tree species
- Reproduces mainly by stump sprout
- Sprouts exhibit rapid initial growth
- Generally slow growing beyond juvenile stage
- Tall and stately on good sites, low and shrubby on poor sites
- May exceed 500 years age in some instances
- Seldom exceed 110 ft height and 60 inches dbh
- Trees 50 to 60 years old often have well developed root burl from which extensive root system spreads
- Requires partial shade for regeneration from seed, and increasing light as tree develops
- Intermediate in shade tolerance
- Persistent foliage

California-Laurel or Myrtlewood (*Umbellularia californica*)

- Broadleaved evergreen with leaves persisting 2 to 6 years
- Broad range of climates ranging from cool and humid to hot and dry
- Tolerates wide range of soil conditions if soil moisture conditions are favorable
- Sprouts prolifically from root collar
- Often multiple-stemmed
- May reach 80 ft in height and 30 inches dbh
- Has been reported as large as 175 ft in height and 159 inches dbh
- Generally slow growing (particularly in height)
- Usually develops deep and spreading root system, but exceptions are common in southern Oregon
- Shade tolerant

Golden Chinkapin or Giant Chinkapin (*Castanopsis chryso-phylla*)

- Broadleaved evergreen
- Often found intermixed in conifer-dominated stands
- Found near sea level in Coast Range to over 5000 ft in Cascade Range, with shrub form found as high as 6000 ft

Golden Chinkapin or Giant Chinkapin (*Castanopsis chryso-phylla*) (continued)

- Climate characterized by winter precipitation and summer drought
- Found in wide variety of topographic position
- Pure stands are uncommon
- Sprouts prolifically when injured, cut, or lightly burned
- Sprouts grow rapidly in height
- Adapted to frequent fire regimes
- May exceed 150 years age
- May reach 150 ft in height and 6 ft dbh
- Tolerates sites with low fertility
- Shrub form is quite tolerant of shade, tree form is less so (intermediate shade tolerance)

References

Burns, Russell M., and Barbara H. Honkala. 1990. *Silvics of North America. Volume 1, Conifers. Volume 2, Hardwoods.* USDA Forest Service. Agricultural Handbook 654.

Jensen, Edward C., and Charles R. Ross. 1994. *Trees to Know in Oregon.* Extension Circular 1450. Oregon State University Extension Service and Oregon Department of Forestry.