



Canola

D.T. Ehrensing

History

Rapeseed and canola are closely related members of the mustard family (Brassicaceae) that are both grown as oilseed crops. All current varieties of rapeseed and canola were developed from *Brassica napus* and *Brassica rapa*.

Rapeseed is grown primarily as a source of erucic acid, which is not edible but is valuable in high-performance industrial lubricants. In the early 1970s, Canadian plant breeders used conventional breeding techniques to remove the anti-nutritional erucic acid and bitter glucosinolates from rapeseed. Removing these compounds resulted in an oilseed crop that produced edible oil low in saturated fats and a very palatable, high-protein meal for animal feed. They coined the word canola (for Canadian oil low acid) to describe a crop that is low in both compounds.

Canada is currently the largest single producer of canola, with nearly 12 million acres grown annually. Because winter canola will not consistently survive their extreme winter conditions, growers on the Canadian prairie produce spring-planted canola.

The Northern Plains of North Dakota and Minnesota currently account for over 90 percent of the 1 million canola acres planted in the United States. Improvements in winter canola varieties in recent years have spurred production in Oklahoma and Kansas, where 60,000 acres were planted in 2006. With continued strong demand for canola oil in food products and as a biodiesel feedstock, canola acreage is expected to increase in all U.S. regions.



Canola flowers and buds.
Photo: Tom Chastain, associate professor, Crop and Soil Science, Oregon State University

Uses

Nutrition

Canola oil is used mainly as cooking oil and in shortening and margarine. To be considered canola, the oil and meal must both meet the following standard:

- Oil < 2% erucic acid
- Meal < 30 micromoles of glucosinolates per gram

The fatty acid profiles of rapeseed and canola are compared in Table 1.

Table 1. Typical fatty acid composition of canola and rapeseed oils.

| Oil source | 16:0 ^a | 18:0 | 18:1 | 18:2 | 18:3 | 20:1 | 22:1 | 22:2 | 24:1 |
|------------|-------------------|------|------|------|------|------|------|------|------|
| Canola | 4 | 2 | 60 | 20 | 10 | — | 2 | — | — |
| Rapeseed | 3 | 1 | 24 | 15 | 8 | 13 | 35 | — | 1 |

^a Fatty acid profiles show the percentage of each fatty acid component in a vegetable oil. The first number in the notation at the top of each column in the profile (e.g., 18:3) indicates the number of carbon atoms in the fatty acid. The number after the colon indicates the number of double bonds in the fatty acid. Although fatty acid profiles vary somewhat from sample to sample, they are generally used to characterize vegetable oils from particular species or varieties of plants.

Because canola oil is very high in unsaturated fatty acids, it is considered a high-quality food oil that is healthy in human diets. Canola oil is particularly desirable for frying because it has a neutral flavor and can be heated to higher temperatures than many other oils without smoking or burning.

Animal feed

Canola meal is a high-protein animal feed used by the dairy, cattle, and poultry industries. It is a byproduct of oilseed extraction that consists of the solids left after oil is extracted from seeds. The meal is highly palatable and typically contains 34 to 38 percent protein with a high percentage of bypass protein. Extracting oil with mechanical screw presses (rather than solvent extraction) usually leaves 8 to 12 percent residual oil in the meal, which also provides a dietary source of energy in animal feed rations.

Industry

Recent petroleum price hikes have greatly increased interest in growing canola for production of biodiesel in the Pacific Northwest (PNW). With its combination of high seed yield and high oil content, canola currently offers the greatest potential oil yield per acre of any PNW crop. High levels of unsaturated fatty acids in canola oil also result



in biodiesel with superior low-temperature performance compared to many other vegetable oils.

Adaptation

Climate

Because it can germinate and grow in relatively cool temperatures, canola is one of the few oilseed crops that can be cultivated over wide areas of the temperate zone. Both winter and spring varieties of canola have been developed, but winter canola normally produces about twice the yield of spring canola in the same production area (see “Yield,” page 8). Canola can also be grown dryland or under irrigation.

A major risk for production of winter canola in the PNW is stand establishment failure. Dry fall weather without enough available soil moisture to germinate and establish non-irrigated winter canola can produce erratic stands or complete crop establishment failure. Late establishment of winter canola can result in significant yield reduction. High temperatures shortly after seedling emergence can also damage and kill young canola plants by burning the stems at the soil surface.

Soil

Canola performs best in well-drained soils and generally will not tolerate flooded or poorly drained areas. Trials are currently underway in the Willamette Valley to determine the performance of winter canola on poorly drained soils that have been tilled to improve drainage.

Cultural practices

Canola fits very well in rotation with many crops, including small grains, grass seed, and potatoes. Due to the potential buildup of diseases, however, do not plant canola more than once every 4 years in a rotation.


Canola is sensitive to many persistent herbicides, such as Glean, Finesse, Assert, atrazine, and triazine compounds. Avoid planting canola in areas where herbicide residues may be a problem. Since canola germinates rapidly, a simple bioassay (planting in soil from the field) can quickly determine if herbicide carryover will affect crop establishment.

Some Brassica species cross-pollinate readily, and isolation of crops is essential. Cross-pollination between edible oil, industrial oil, condiment mustard, and Brassica vegetable seed crops generally reduces quality of all crops. Most PNW states have established Brassica production districts to minimize production conflicts among growers (see “Limitations on canola production in Oregon,” page 8).



Seedbed preparation

Canola seeds are small and require a firm seedbed free of large clods for optimum germination and stand establishment. Most growers use a roller with the last tillage operation to pack the soil before planting. Working soil too fine or packing too hard can cause crusting, which may inhibit seedling emergence.



Canola seeds.

Planting date

Since potential yield declines rapidly with later fall planting, optimum seeding dates are especially important when growing winter canola. The goal is to grow well-established plants with a closed canopy of leaves before the onset of cold weather. In western Oregon, winter canola typically is planted in mid-September. In eastern Oregon, begin planting winter canola in mid-August. Plant spring canola at the normal time for spring grain crops.

Seeding rate

Suggested seeding rates for winter canola range from 4 to 12 pounds of seed per acre. Under normal conditions, plant 4 to 7 pounds of seed per acre. Use higher seeding rates (8 to 12 pounds per acre) for late plantings, heavier soils, or in field conditions where emergence may be reduced. Plant spring canola at 5 to 8 pounds of seed per acre.



Young canola plant.

Photo: Tom Chastain, associate professor, Crop and Soil Science, Oregon State University

Seeding depth

Optimum seeding depth is $\frac{1}{2}$ to 1 inch.

Row spacing

Canola usually is planted with a conventional grain drill. In high-moisture areas or under irrigation, a row width of 6 to 10 inches normally is used. In dryland conditions, row width should be increased to 12 to 16 inches. Narrower row spacing promotes rapid canopy closure and reduces weed competition.

Variety selection

Many commercial winter and spring canola varieties are available. Most of the recently developed spring canola varieties have been



genetically modified to induce herbicide resistance. These include 'Clearfield', 'Liberty Link', and 'Roundup Ready'. Herbicide resistant winter canola varieties are also beginning to appear on the market. Non-GMO winter and spring canola varieties are available, but they are rapidly becoming difficult to find.

Fertilizer

Response of canola to fertilizer application is highly dependent on moisture availability and soil type. Under irrigation or in high rainfall areas where yield potential is higher, increased rates of applied fertilizer may be economical, while less may be needed for maximum production in dryland conditions or on less productive soils.



Canola rosettes at 10 weeks.

Nitrogen

Nitrogen is generally the most limiting nutrient in canola production, so providing adequate nitrogen is essential to maximize canola yield. The amount of nitrogen fertilizer required by canola depends on the variety, its potential yield, and the amount of available nitrogen in the soil. Canola takes up approximately 7 lb N/100 lb of expected seed yield. If fall N is needed for winter canola, apply 30 to 50 lb N/acre prior to planting and the remainder in the spring.

Since most nitrogen uptake in canola occurs before bolting, adequate N must be available in the soil prior to the crop's rapid growth period. Apply spring N before bolting begins.

Sulfur

Sulfur requirements of canola are greater than those of cereal crops. Canola requires about 1 lb S/100 lb of expected seed yield.

Phosphorus

Both winter and spring canola often benefit from application of phosphorus when soil test levels are below 5 ppm. Phosphorus is generally most effective when applied prior to seeding.

Potassium

Soil potassium levels normally are adequate for canola in the PNW, but apply potassium when soil test values are below 75 ppm.

Boron


Winter canola generally requires more boron than many other crops grown in rotation with it. Apply 1 to 2 lb boron/acre if soil tests indicate levels below 0.5 ppm. Avoid over-application, since excess boron is toxic to canola.



Weed control

Canola seedlings normally develop quickly and compete well with annual weeds. Well-timed planting and good stand establishment promote rapid leaf canopy closure, which suppresses many weed species. Weed problems usually develop with late planting and poor stand establishment.

Several grass and broadleaf herbicides are currently registered for use in canola production in the PNW. Consult local weed control guides for information on application rates and timing.



Canola plants in bloom.

Pest management

Diseases

Plant diseases are one of the most serious canola production problems. The two most important diseases are sclerotinia stem rot and blackleg.

Sclerotinia stem rot (or white mold) is caused by the fungus *Sclerotinia sclerotiorum*. Infection weakens the plant stem, causing losses from lodging and early ripening. This disease infects many other crops, including sunflower, potatoes, safflower, beans, peas, and alfalfa.

Blackleg, the most serious disease of canola, is caused by the fungus *Phoma lingam* (*Leptosphaeria maculans*). This disease is carried over from season to season on infected stubble and spreads by airborne spores or through infected seed. Seed treatment with fungicides is very effective in controlling blackleg and is required on planting seed used in Oregon. Blackleg can cause yield reductions up to 50 percent due to premature ripening and lodging. To date, blackleg has not been found in the PNW.

Since no fungicides are currently available for use on canola, it's important to practice a minimum 4-year crop rotation. Use of certified, treated seed is required for canola planting in Oregon. Using resistant varieties will also help reduce disease problems as canola production increases.

Insect pests

Insects have been relatively minor pests in PNW canola production, but several insect pests have been identified in canola fields in Oregon.

Cabbage seedpod weevil (*Ceutorynchus assimilis*) is the most important insect pest of canola. Adult weevils lay eggs near young seedpods, and the larvae enter the pods to feed on developing seeds. Larval feeding can reduce yield by 20 to 30 percent. Field monitoring and proper timing of insecticide treatment are essential to achieve control.



Several species of aphids (cabbage aphid, turnip aphid, and green peach aphid) are found on canola in the PNW. Aphid feeding on rosettes can reduce rosette size and vigor, adversely affecting winter survival and yield. Aphid feeding on floral parts can reduce seed set and yield.

Adult flea beetles sometimes damage canola seedlings by feeding on cotyledons and young leaves, causing a shothole appearance. This usually occurs during warm, dry weather conditions shortly after crop emergence.

Harvest

Proper harvest timing is critical with canola to attain high yield and oil quality. Early harvest leads to excessive green seed, reduced oil content, and high seed moisture. Late harvest can result in severe crop losses due to shattering, because mature, dry canola pods open very easily.

Harvest canola either by swathing followed by combine pickup when the crop is dry, or by direct combining. Timing of swathing is important to minimize seed losses. Because canola seeds on the bottom of the plant mature before those at the top, sample seed from pods on the lower third of the main stem when inspecting the crop for maturity. Begin swathing when seeds have turned brown on the lower third of the main stem.



Windrowing canola.



Canola pods.

Photo: Tom Chastain, associate professor, Crop and Soil Science, Oregon State University



Combining canola.



Oilseed Crops

Direct combining standing canola can be extremely risky, and seed losses can be very high when the pods are dry and brittle. In the Columbia Basin, growers have had good success using “canola pushers” prior to combine harvesting. The pushers bend the canola stems over just above the soil surface rather than cutting the stems with a windrower. This allows the crop to ripen while reducing stem movement and shattering by the wind. When mature, the canola is then direct combined traveling in the opposite direction of the pusher.

Fully mature canola seed is dark brown or black. Optimum seed moisture at harvest is 8 to 9 percent.

Yield

Canola yield varies widely depending on variety, time of planting, available moisture, soil fertility, and harvest timing. Well-established winter canola tends to produce the highest seed yield.

In recent variety trials in the Willamette Valley, winter canola yield ranged from approximately 1,900 to over 4,800 pounds of seed per acre. Spring canola tends to produce about half the yield of winter canola in the Willamette Valley.

Limitations on canola production in Oregon

In the early 1990s, the Oregon legislature gave the director of the Oregon Department of Agriculture (ODA) authority to regulate the production of rapeseed (plants in the genus *Brassica*). Brassicaceae is a large and widespread plant family including broccoli, cauliflower, cabbage, turnips, radishes, rapeseed, canola, mustard, and others. The purpose of this law was to minimize cross-pollination conflicts among vegetable seed growers and seed companies producing these important crops.

In 1993, Oregon Administrative Rules (Chapter 603, Division 52) were adopted establishing rapeseed production districts and isolation requirements for producers. A system of pinning maps was also established in the Linn and Marion county Extension offices to facilitate maintenance of isolation distance between vegetable seed fields. Under this system, growers or seed companies mark out areas on a map with pins to identify vegetable production fields on a first-come, first-served basis. Producers attempt to maintain a 3-mile isolation distance between Brassica crops that are probable cross-pollinators.

Because of increased fuel prices and canola’s excellent adaptation to the PNW, grower interest in producing canola for biodiesel has greatly increased in the last several years. In response to this interest and concerns from Willamette Valley vegetable seed producers, the ODA



revised the Brassica rules in 2005, severely restricting areas in the state where canola can be grown as an oilseed crop. Districts where vegetable seed production is protected are shown on the map in Figure 1.




Figure. 1. Protected vegetable seed production districts under the 2005 canola rules.

Within the protected districts, growing any Brassica species for oil production is prohibited. Under current rules, producers wishing to grow Brassicas for oil within a protected district may request an exemption from the director of the ODA.

Several applications for exemption were submitted to the director in 2006. Exemptions were denied for Willamette Valley and Central Oregon commercial growers, but a permit was issued to allow research on the Central Oregon agricultural experiment stations provided that no genetically modified canola was included in the trials. In fall 2006, an exemption was granted to allow Oregon State University to conduct limited canola trials in the Willamette Valley as part of a program to assess the commercial potential of a number of oilseed crops in the area.

The rules were reviewed and maintained without change in fall 2007. Current rules will be open for review and possible modification in 2009.

Bibliography

- Berglund, D.R. and Kent McKay. 2002. Canola Production. North Dakota State University Extension Circular A-686.
<http://www.ag.ndsu.edu/pubs/plantsci/crops/a686w.htm>
- Franzen, D.W. 1999. Fertilizing Mustard and Canola. North Dakota State University Extension Circular SF-1122.
<http://www.ag.ndsu.edu/pubs/plantsci/soilfert/sf1122w.htm>
- Oregon Department of Agriculture. 2007. Rapeseed Control Areas.
http://www.oregon.gov/ODA/PLANT/canola_rapeseed.shtml

(Continued next page)



Wysocki, D.J., M. Corp, D.A. Horneck, and L.K. Lutcher. 2007. Irrigated and Dryland Canola Nutrient Management Guide. Oregon State University, EM 8943-E.

<http://extension.oregonstate.edu/catalog/pdf/em/em8943-e.pdf>

Zollinger, R.K. 2007. 2007 North Dakota Weed Control Guide. North Dakota State University Extension Circular W-253.

<http://www.ag.ndsu.edu/weeds/w253/w253-1d.htm>

Trade-name products and services are mentioned as illustrations only. This does not mean that the Oregon State University Extension Service either endorses these products and services or intends to discriminate against products and services not mentioned.

© 2008 Oregon State University

This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties.

Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

Published February 2008.

