

# Storing Pumpkin and Winter Squash at Home

N.S. Mansour

All pumpkins and hard-shelled winter squash may be stored at the end of the growing season for use well into the new year. For best results, store sound, well-cured fruit at 50 to 55°F in a 50 to 70% relative humidity.

Length of storage life varies according to variety and type of squash or pumpkin.

## *Hard-shelled winter squash*

	<i>Storage life</i>
Table Queen (acorn type)	1 to 2 months
Butternut	2 to 3 months
Hubbard types	3 to 6 months
Banana	3 to 6 months
Buttercup (turban type)	3 to 6 months
Sweet Meat	4 to 6 months

## *Pumpkin*

Jack O'Lantern	2 to 3 months
Connecticut Field	2 to 3 months

Table Queen and other acorn-type squash can be stored satisfactorily for 1 to 2 months. With longer storage, the skin begins to turn yellow and the squash becomes stringy.

## **Harvesting**

Harvest all types of squash and pumpkin before frost begins. Squash are ready for harvest when the rind is hard enough to resist fingernail scratches. Cut the stem 2 to 4 inches from the fruit. Pumpkins without stems do not store well. Hubbard-type squash stores best with the stems completely removed. Handle fruit carefully to keep them in good condition. Slightly immature squash and pumpkin can be used if they are cured properly. Curing helps toughen the skins of immature fruit and helps heal cuts and scratches.

## **Curing**

All squash undergo a slow curing process during proper storage. Artificial curing is not necessary for well matured squash under good storage conditions. Nearly mature squash, except acorn types, may benefit from a short period of curing. Curing is holding squash and pumpkin at a temperature favorable for healing cuts and scratches and for forming a protective corky layer over injuries and cut surfaces of the stem.

Cure squash and pumpkin for 10 days at temperatures of 80 to 85°F and a relative humidity of 80 to 85 degrees. Use a small cabinet heated by a thermostatically controlled electric heater or a corner of the garage partitioned off with plastic for a curing chamber. A small fan will maintain good circulation and uniform distribution of heat.



## **Storing**

Squash and pumpkin deteriorate rapidly if stored at temperatures below 50°F. The best storage temperature is between 50 and 55°F. Fruit that has been exposed to freezing before harvest also will deteriorate rapidly. A relative humidity of 75%, about normal for garages or other suitable storage areas in western Oregon, is satisfactory. Keep the temperature at 50 to 55°F. Keep the surface of the fruit dry to prevent or retard growth of decay fungi and bacteria. Air circulation helps to prevent moisture from forming on the surfaces of the fruit.

Provide shelves for storage of pumpkin and squash. Do not store fruit on cold concrete floors. Promptly discard any fruit that shows signs of decay. Some of the more durable squash may be stacked on top of each other if adequate room is provided for air circulation. Do not store pumpkin or squash near apples, pears, or other ripening fruit. Ripening fruit releases ethylene gas, which causes yellowing of the squash and shortens storage life.

*N.S. Mansour, Extension vegetable crops specialist, Oregon State University.*



Extension Service, Oregon State University, Corvallis, O.E. Smith, director. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties.

Oregon State University Extension Service offers educational programs, activities, and materials—without regard to race, color, national origin, sex, age, or disability—as required by Title VI of the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and Section 504 of the Rehabilitation Act of 1973. Oregon State University Extension Service is an Equal Opportunity Employer.



**OREGON STATE UNIVERSITY EXTENSION SERVICE**