Spotted Wing Drosophila (SWD)

A quick, 7-step guide for detecting SWD larvae in fruit

EM 9097 • October 2014

Amy J. Dreves, Adam Cave, and Jana Lee

STEP 1: Gather supplies



- ☐ White or black shallow tray
- ☐ Magnifying lens or magnifying headset, such as an OptiVisor
- ☐ Sealable plastic bags
- ☐ Forceps, fine paintbrush
- ☐ Stirring stick or spoon

☐ Salt or light brown sugar and water

Extension

Service

Drosophila suzukii

- ☐ 1-gallon jug to hold solution
- ☐ Good lighting
- Optional: A wire mesh screen to submerge fruit

STEP 2: Prepare 1-gallon solution



- ☐ At least one day before testing, dissolve 1 cup of plain salt or 2.5 cups of light brown sugar in 1-gallon jug of warm water.
- ☐ Agitate solution jug to hasten dissolving
- ☐ Re-agitate the solution the day of testing

STEP 3: Collect fruit



- ☐ Collect fruit with symptoms of damage from SWD or collect random samples of fruit
- ☐ Place fruit in a sealable plastic bag
- ☐ The more fruit you inspect, the more confidence you will have in the quality of the rest of the fruit.
- ☐ Label bag to reference collection location

(For example: 8/2/14, row 88, plant #5, NE side)

Amy J. Dreves, assistant professor-senior research and Extension entomologist, Crop and Soil Science Department, Oregon State University; Adam Cave, biological science technician; and Jana Lee, research entomologist, U.S. Department of Agriculture-Agricultural Research Service, Horticulture Crops, Corvallis, OR.

STEP 4: Crush fruit in bag



- ☐ Crush fruit, but do not make a purée
- ☐ A rolling pin makes it easy to mash up the fruit and expose the pulp
- Use your fingers to crush any fruit not opened
- ☐ Avoid aggressive crushing that will damage larvae

STEP 5: Add solution to bag



- ☐ Mix and allow solution to penetrate fruit pulp
- ☐ Swish solution to distribute it among fruit debris
- ☐ Solution will irritate any larvae inside fruit and encourage them to leave fruit and rise to surface

STEP 6: Pour contents into shallow tray



- ☐ Cover crushed fruit completely with salt or sugar solution
- ☐ Most larvae will float to top for easy viewing
- ☐ Sugar will keep SWD alive longer, to see larvae move or to raise larvae to confirm species of fly

STEP 7: Examine for SWD larvae





- ☐ Choose a well-lit area
- Use magnifier (hand lens) or headset magnifier
- Examine surface closely for bobbing or floating larvae (approximately 0.7mm–5.4mm long)
- ☐ Look for larvae in the first few to 15 minutes
- ☐ Stir contents periodically to dislodge larvae from fruit and tray sides
- ☐ Use a wire screen to submerge fruit and to allow larvae to float
- ☐ Do not mistake SWD for plant parts, aphid skins, fruitworms, thrips, pulp threads, sap beetle larvae, larvae from other *Drosophila* species
- ☐ Count and record larvae, and track seasonal changes in numbers
- ☐ Use tab counter to record number of larvae
- ☐ The 7-step process described here is a guide and not a guarantee that you do not have an infestation somewhere in your crop.

Project supported by Agriculture and Food Research Initiative Competitive Grant No.: 2010-51181-21167; USDA National Institute of Food and Agriculture, Oregon State University-Crop and Soil Science Department, and U.S. Department of Agriculture, Agricultural Research Station, Horticulture Crops.

Trade-name products and services are mentioned as illustrations only. This does not mean that the Oregon State University Extension Service either endorses these products and services or intends to discriminate against products and services not mentioned.

© 2014 Oregon State University. This publication was produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.