

Optimizing chemical control of spotted-wing drosophila

Serhan Mermer, Gwen-Alyn Hoheisel, Haitham Bahlol, Lav Khot, Dalila Rendon, Linda Brewer, Daniel Dalton, Rachael Nieri, Kyoo Park, Ferdinand Pfab, Marco Rossi-Stacconi and Vaughn Walton

Key points

- Most spotted-wing drosophila are found in the center of the fruit crop canopy, where they find shade, cooler temperatures and higher humidity.
- Sprayer calibration, nozzle orientation, and adequate spray volume are key to good coverage.
- Apply sprays using appropriate tractor speeds.
- Air-blast sprayers provide consistent spray coverage.
- Cannon sprayers minimize disturbance of fruit close to harvest and cover large areas rapidly.
- Electrostatic sprayers minimize application time, non-target deposition and fruit disturbance while providing adequate coverage.
- No sprayer type provides complete coverage.




Figure 1. Advantages and disadvantages of three types of spray equipment used in small fruit production.

Credit: Marco Rossi-Stacconi, © Oregon State University

Spray equipment and crop canopy are key factors

Crop canopy and structure and the type of pesticide spray equipment help determine the effectiveness of pesticide applications. Spotted-wing drosophila lay eggs in the center of the canopy rather than in the upper and outer portions. Thick canopies impede the penetration of insecticides into this habitat, resulting in reduced control. Plants

that are more heavily and appropriately pruned allow improved targeting of the interior of the canopy with the pesticide and improve pest control.

Pest control also depends on spray equipment settings and maintenance. Tractor speed, nozzle type, pressure and spray volume all affect insecticide spray efficacy. Evaluate these factors from time to time according to the size and structure of the crop. Three common types of sprayers are used in Oregon small fruit crop production: air-blast, cannon and electrostatic sprayers (Figure 1).

Cannon sprayers reduce application time and require lower spray volumes than air-blast sprayers. When used for border sprays, cannon sprayers project the chemical across the top of several crop rows from the outside of the field without covering the whole block.

Border spray programs using cannon sprayers result in similar levels of SWD control in Oregon blueberry compared to complete block applications while reducing impact to nontarget arthropods.

However, cannon sprayers may result in irregular spray deposition and increased off-target drift under windy conditions. Irregular deposition patterns may reduce control when SWD pressure is high.

Air-blast sprayers rely on rapid airflow for more uniform deposition of pesticide on all surfaces throughout the canopy. In recent tests, air-blast sprayers increased spray deposition on Oregon blueberry on the lower part of the canopy. This could be because the axial fan circulates the air from the bottom to the top of the canopy. Orienting the nozzle to provide adequate coverage optimizes the deposition pattern within the canopy. Nozzles should be checked and serviced annually.

Electrostatic sprayers deliver pesticides to hard-to-reach locations within the canopy. The electrically charged droplets are drawn to the oppositely charged plants. This reduces off-target product losses to the air and ground and delivers more uniform canopy coverage. It is possible for high humidity to negatively affect coverage uniformity and adequacy. This equipment will likely cost more than air-blast and cannon sprayers.

Resources

- Diepenbrock, L. M., H.J. Burrack. 2016. Variation of within-crop microhabitat use by *Drosophila suzukii* (Diptera: Drosophilidae) in blackberry. *Journal of Applied Entomology*. 141: 1–7.
- Klick, J., W.Q. Yang, J.C. Lee, D.J. Bruck. 2016. Reduced spray programs for *Drosophila suzukii* management in berry crops. *International Journal of Pest Management*. 62(4): 368–377.
- Khot, L.R., R. Ehsani, G. Albrigo, P.A. Larbi, A. Landers, J. Campoy, C. Wellington. 2012. Air-assisted sprayer adapted for precision horticulture: Spray patterns and deposition assessments in small-sized citrus canopies. *Biosystems Engineering*. 113, 76–85.
- Singh, M., C. Ghanshyam, P.K. Mishra, R. Chak. 2013. Current status of electrostatic spraying technology for efficient crop protection. *AMA, Agricultural Mechanization in Asia, Africa and Latin America*. 44: 46–53.
- VanEe, G., L.R. Ledebuhr, E. Hanson, J. Hancock, D.C. Ramsdell. 2000. Canopy development and spray deposition in highbush blueberry. *HortTechnology* 10: 353–359.
- Mermer, S., F. Pfab, G.A. Hoheisel, H.Y. Bahlol, L. Khot, D.T. Dalton, L.J. Brewer, M.V. Rossi-Stacconi, C. Zhang, L. Xue, V.M. Walton. [Canopy spray deposition and related mortality impacts of commonly used insecticides on *Drosophila suzukii* Matsumura \(Diptera: Drosophilidae\) populations in blueberry. \(https://doi.org/10.1002/ps.5672\)](https://doi.org/10.1002/ps.5672)

About this series

This publication is one of a series of nine focused on strategies for controlling spotted-wing drosophila in Oregon. Learn more:

- EM 9261: [How seasons affect population structure, behavior and risk on spotted-wing drosophila](https://catalog.extension.oregonstate.edu/em9261) (<https://catalog.extension.oregonstate.edu/em9261>)
- EM 9262: [Cultural control strategies to manage spotted-wing drosophila](https://catalog.extension.oregonstate.edu/em9262) (<https://catalog.extension.oregonstate.edu/em9262>)
- EM 9263: [Host range and characteristics affecting fruit susceptibility to spotted-wing srosophila](https://catalog.extension.oregonstate.edu/em9263) (<https://catalog.extension.oregonstate.edu/em9263>)
- EM 9264 [Noncrop host plants used by spotted-wing drosophila](https://catalog.extension.oregonstate.edu/em9264) (<https://catalog.extension.oregonstate.edu/em9264>)
- EM 9265: [Chemical control of spotted-wing drosophila: Spray applications](https://catalog.extension.oregonstate.edu/em9265) (<https://catalog.extension.oregonstate.edu/em9265>)
- EM 9266: [Chemical control of spotted-wing drosophila: Insecticide efficacy](https://catalog.extension.oregonstate.edu/em9266) (<https://catalog.extension.oregonstate.edu/em9266>)
- EM 9267: [Monitoring techniques for spotted-wing drosophila](https://catalog.extension.oregonstate.edu/em9267) (<https://catalog.extension.oregonstate.edu/em9267>)
- EM 9268: [Potential impacts of irrigation on biocontrol on spotted-wing drosophila populations](https://catalog.extension.oregonstate.edu/em9268) (<https://catalog.extension.oregonstate.edu/em9268>)
- EM 9269: [Biocontrol of spotted-wing drosophila](https://catalog.extension.oregonstate.edu/em9269) (<https://catalog.extension.oregonstate.edu/em9269>)

About the authors

Serhan Mermer

Graduate student, environmental and molecular toxicology
Department of Horticulture, Oregon State University

Gwen-Alyn Hoheisel

Regional Extension Specialist
Oregon State University

Haitham Bahlol

Graduate student
Department of Biological Systems Engineering, Washington State University

Lav Khot

Assistant professor

Department of Biological Systems Engineering, Washington State University



[Dalila Rendon \(https://extension.oregonstate.edu/people/dalila-rendon\)](https://extension.oregonstate.edu/people/dalila-rendon)

Assistant Professor (Practice), Department of Horticulture



[Linda Brewer \(https://horticulture.oregonstate.edu/users/linda-brewer\)](https://horticulture.oregonstate.edu/users/linda-brewer)

Senior Faculty Research Assistant II, Department of Horticulture

Daniel Dalton

Graduate student, horticultural entomology

Department of horticulture, Oregon State University

Rachael Nieri

Post-doctoral researcher, horticultural entomology

Department of Horticulture, Oregon State University

Kyoo Park

Graduate student

Department of Horticulture, Oregon State University

Ferdinand Pfab

Post-doctoral researcher, ecology, evolution and marine biology

University of California, Santa Barbara

Marco Rossi-Stacconi

Post-doctoral researcher, horticultural entomology
Department of Horticulture, Oregon State University



Vaughn Walton (<https://extension.oregonstate.edu/people/vaughn-walton>)

Entomologist

Related publications



Credit: © Oregon State University (Cropped from original)

Biocontrol of spotted-wing drosophila

(<https://extension.oregonstate.edu/catalog/em-9269-biocontrol-spotted-wing-drosophila>)

Biological control agents target SWD, predators, parasitoids, nematodes and microorganisms such as fungi and bacteria. Predators eat SWD, removing a substantial portion from the production system.

Marco Rossi-Stacconi, Linda Brewer, Elizabeth Miller, Daniel Dalton, Jana Lee, Kyoo Park, Ferdinand Pfab, Vaughn Walton, Cherre Bezerra Da Silva | Sep 2024 | OSU EXTENSION CATALOG [Peer reviewed \(Orange level\)](#) (<https://extension.oregonstate.edu/peer-review-guidelines>)



Potential impacts of irrigation on biocontrol of spotted-wing drosophila

(<https://extension.oregonstate.edu/catalog/pub/em-9268-potential-impacts-irrigation-biocontrol-spotted-wing-drosophila>)

A naturally occurring parasitic wasp in Oregon can help control spotted-wing drosophila in blueberries, but overhead irrigation can limit the effectiveness of this biocontrol method.

Cherre Bezerra Da Silva, Brianna F. Price, Daniel Dalton, Dalila Rendon, Kyoo Park, Linda Brewer, Vaughn Walton, Marco Rossi-Stacconi | Sep 2024 | OSU EXTENSION CATALOG [Peer reviewed \(Orange level\)](#) (<https://extension.oregonstate.edu/peer-review-guidelines>)



Improved chemical control strategies for spotted-wing drosophila

<https://extension.oregonstate.edu/catalog/em-9266-improved-chemical-control-strategies-spotted-wing-drosophila>

Learn how life stages, product choice, timing and coverage impact success in the control of spotted-wing drosophila.

Serhan Mermer, Linda Brewer, Daniel Dalton, Rachael Nieri, Kyoo Park, Ferdinand Pfab, Marco Rossi-Stacconi, Vaughn Walton | Sep 2024 |

OSU EXTENSION CATALOG [Peer reviewed \(Orange level\)](#)

<https://extension.oregonstate.edu/peer-review-guidelines>

© 2019 Oregon State University. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination on the basis of race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, familial/parental status, income derived from a public assistance program, political beliefs, genetic information, veteran's status, reprisal or retaliation for prior civil rights activity. (Not all prohibited bases apply to all programs.)

Accessibility: This publication will be made available in an accessible alternative format upon request. Please contact puborders@oregonstate.edu or 541-737-3311.